

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

J. Differential Equations 226 (2006) 525–540

**Journal of
Differential
Equations**

www.elsevier.com/locate/jde

An existence result for a class of second order evolution equations of mixed type

Fabio Paronetto

Dipartimento di Matematica “Ennio De Giorgi”, Università degli Studi di Lecce, Via per Arnesano, 73100 Lecce, Italy

Received 11 April 2005; revised 12 December 2005

Available online 9 February 2006

Abstract

We give an existence and uniqueness result for a linear abstract evolution equation of second order with some coefficient in front of the second temporal derivative which may degenerate to zero and change sign.
© 2006 Elsevier Inc. All rights reserved.

Keywords: Differential equations of mixed type; Second order evolution equations

1. Introduction

We present an existence result for the solution of the abstract evolution equation of second order

$$(Su')' + \mathcal{N}u' + \mathcal{A}u = f \quad (1)$$

with suitable data. Given V Banach space and H Hilbert space, $V \subset H$, defined $\mathcal{V} = L^2(0, T; V)$, \mathcal{S} will be a linear operator defined in $L^2(0, T; H)$ and the solution u will be taken in $C([0, T]; V)$, with $u' \in \mathcal{V}$ and $(Su')' \in \mathcal{V}'$. Notations and definitions for the abstract equation will be given in the second section, so for the moment we confine ourselves to explain a concrete and simple problem (see also Section 3 and [8]): suppose to have a bounded open set Ω of \mathbf{R}^n , $T > 0$, a function $s : \Omega \times [0, T] \rightarrow \mathbf{R}$, $s \in L^\infty(\Omega \times (0, T))$, define $\Omega_+(t) = \{x \in \Omega \mid s(x, t) > 0\}$ and $\Omega_-(t) = \{x \in \Omega \mid s(x, t) < 0\}$, consider $H = L^2(\Omega)$ and $V = H_0^1(\Omega)$, and define $Su(x, t) = s(x, t)u(x, t)$. Consider

E-mail address: fabio.paronetto@unile.it.

$$\begin{cases} (s(x, t)u_t(x, t))_t - \Delta u_t(x, t) - \Delta u(x, t) = f(x, t) & \text{on } \Omega \times (0, T), \\ u(x, t) = 0, & (x, t) \in \partial\Omega \times (0, T), \\ u_t(x, 0) = \varphi(x), & x \in \Omega_+(0), \\ u_t(x, T) = \psi(x), & x \in \Omega_-(T), \\ u(x, 0) = \eta(x), & x \in \Omega, \end{cases} \quad (2)$$

where $f: \Omega \times (0, T) \rightarrow \mathbf{R}$, $\varphi: \Omega_+(0) \rightarrow \mathbf{R}$, $\psi: \Omega_-(T) \rightarrow \mathbf{R}$, $\eta: \Omega \rightarrow \mathbf{R}$ are the data of the problem. Notice that for this kind of problem we prescribe an initial datum for the function u in the whole Ω , but for the function u_t only in the part in which at time $t = 0$ the coefficient s is positive, while at time $t = T$ where s is negative. Where $s \equiv 0$ no data for u_t are imposed. These requirements about initial/final data for u_t are quite natural and some comments about them will be done in Section 3.

Equation like that in (2) are widely studied. The classical case corresponds to take $s \equiv 1$ (or s strictly positive) in (2), i.e. the wave equation with dissipation or friction (see, for instance, [16, Section 33.5]).

Also the theory of elasticity (see, for instance, [4, Chapter 3]) leads to an equation like

$$u_{tt} = \operatorname{div}(a(Du)),$$

in which often an additional dissipative term is added (a term $\epsilon \Delta u_t$ with ϵ positive and little parameter) to regularise the solution, so that the equation becomes (see, e.g., [10,11,15] and, for a non-monotone, for instance, [9])

$$u_{tt} = \operatorname{div}(a(Du) + \epsilon Du_t). \quad (3)$$

Moreover the quasi-steady approximation, which correspond to consider $s \equiv 0$, i.e.

$$\operatorname{div}(a(Du) + \epsilon Du_t) = 0, \quad (4)$$

has been also considered for equation of this type (we refer, e.g., for this case, to [12]).

Changing type equations have been already studied a lot. As regards evolution equations of first order (partially elliptic, partially parabolic) we recall some classical books (see [2,14]) where many degenerate problems are presented in which the coefficients in front of temporal derivatives may be non-negative. Among the situations in which the coefficient in front of temporal derivative may be also negative we recall [1,7,8].

As regards the degenerate situation for second order equations like that in (2) we recall [13, see in particular Chapter VI] for $s \geq 0$ (one can think there are two regions, one where the evolution is standard like in (3) and a region where the evolution is quasi-steady like in (4)).

A well-known and studied situation of changing type equation of second order is the Tricomi equation

$$xu_{tt} - u_{xx} = 0,$$

or more generally $s(x)u_{tt} - u_{xx} = 0$, which occurs when studying transonic flow, for which we refer to the recent and also historical paper [6] (see also the references therein) and to [3].

In the present paper we analyse the situation considered by Showalter in [13], generalising (see Theorem 3.4) the existence result in the sense that the coefficient s may be not only non-negative, but positive, null and negative.

To do this, we reduce by a change of variable to a first order evolution equation and use an existence result for mixed equation of first order (see [8]) which follows by a classical result for perturbation of monotone operators (Theorem 2.5). Although in [8] the operator \mathcal{A} in (1) may be non-linear, here we manage to consider \mathcal{A} only linear (this is due to Proposition 2.3, needed when we reduce to a first order equation).

With our method we do not manage to generalise the Tricomi equation (which correspond to take $\mathcal{N} = 0$ in (1)), neither via regularisation, i.e. taking the solutions of $(Su')' + \epsilon \mathcal{N}u' + \mathcal{A}u = f$ and then let ϵ go to zero (see, for instance, [5, Chapter 3, Section 8.5] for the analogous classical result) due to the change of sign of \mathcal{S} .

At the end we comment a concrete example, where the operator \mathcal{S} is defined by a function s like in (2), referring to [8] for examples of possible choices of the function s .

2. Recalls and notations

We want to give an existence result for the abstract equation

$$(Su')' + \mathcal{N}u' + \mathcal{A}u = f$$

($\mathcal{S}, \mathcal{N}, \mathcal{A}$ are operators) with suitable “initial/final” conditions. In this section we give some recalls, assumptions about these three operators and some results contained in [8].

Consider a triplet

$$V \subset H \subset V'$$

where V is a reflexive and separable Banach space, V' its dual space, H a Hilbert space and the embedding $V \subset H$ is continuous and dense. Fix $T > 0$ and denote

$$\mathcal{V} = L^2(0, T; V), \quad \mathcal{H} = L^2(0, T; H), \quad \mathcal{V}' = L^2(0, T; V').$$

Consider a family of linear self-adjoint operators

$$S(t) : H \rightarrow H, \quad t \in [0, T].$$

For every $t \in [0, T]$ consider the decomposition

$$H = H_+(t) \oplus H_0(t) \oplus H_-(t)$$

where $H_0(t)$ is the kernel of $S(t)$, $H_+(t)$ and $H_-(t)$ are defined respectively as the subspaces of H such that $(S(t)u, u)_H > 0$ for every $u \in H_+(t)$, $u \neq 0$, and $(S(t)u, u)_H < 0$ for every $u \in H_-(t)$, $u \neq 0$, and decompose $S(t)$ as

$$S(t) = S_+(t) + S_0(t) - S_-(t) \quad \text{where}$$

$$S_+(t) : H_+(t) \rightarrow H_+(t) \quad \text{defined by} \quad S_+(t)u = S(t)u \quad \text{for } u \in H_+(t),$$

$$S_-(t) : H_-(t) \rightarrow H_-(t) \quad \text{defined by} \quad S_-(t)u = -S(t)u \quad \text{for } u \in H_-(t) \quad (5)$$

and $S_0(t) : H_0(t) \rightarrow H_0(t)$ is the operator $S_0(t)u = 0$ for every $u \in H_0(t)$. Moreover we define

$$|S(t)| : H \rightarrow H \quad \text{by} \quad |S(t)|u = S_+(t)u + S_-(t)u.$$

Now denote

$$\tilde{H}(t), \tilde{H}_+(t), \tilde{H}_-(t) \quad \text{the completion respectively of} \quad H, H_+(t), H_-(t) \quad (6)$$

with respect to the norm $\|w\|_{\tilde{H}(t)} = \| |S(t)|^{1/2} w \|_H$ and denote the orthogonal projections

$$P_+(t) : \tilde{H}(t) \rightarrow \tilde{H}_+(t), \quad P_-(t) : \tilde{H}(t) \rightarrow \tilde{H}_-(t), \quad P_0(t) : \tilde{H}(t) \rightarrow H_0(t)$$

for every $t \in [0, T]$ and in particular we will need

$$P_+(0) : \tilde{H}(0) \rightarrow \tilde{H}_+(0) \quad \text{and} \quad P_-(T) : \tilde{H}(T) \rightarrow \tilde{H}_-(T). \quad (7)$$

Before giving assumptions about S we recall a definition.

Definition 2.1. We say that $B : [0, T] \rightarrow \mathcal{L}(X, X')$, X Banach space, is differentiable if, for every $u, v \in X$ the function

$$t \mapsto \langle B(t)u, v \rangle_{X' \times X}$$

is absolutely continuous on $[0, T]$ and there exists a function $b \in L^1(0, T)$ such that

$$\left| \frac{d}{dt} \langle B(t)u, v \rangle_{X' \times X} \right| \leq b(t) \|u\|_X \|v\|_X \quad \text{for a.e. } t \in [0, T].$$

Observe that $B'(t) : X \rightarrow X'$ is linear for almost every $t \in (0, T)$.

Remark 2.2. If B is differentiable in the sense just defined, the following formula holds: if $u \in W^{1,p}(0, T; X)$ for any p , then

$$(B(t)u(t))' = B'(t)u(t) + B(t)u'(t) \quad \text{for a.e. } t \in (0, T).$$

In the special case $B(t) = B$ for every t , taking $v \in L^p(0, T; X)$ and $u(t) = \int_0^t v(s) ds$, we also derive from the formula above that

$$B \int_0^t v(s) ds = \int_0^t Bv(s) ds. \quad (8)$$

Assumptions about \mathcal{S}

We suppose that the family of operators $S(t)$ will satisfy:

- (i) $S(t)$ self-adjoint,
- (ii) $\max_{t \in [0, T]} \|S(t)\|_{\mathcal{L}(H)} \leq C_1$,
- (iii) $t \mapsto (S(t)u, v)_H$ absolutely continuous on $[0, T]$,
- (iv) $\Gamma \|u\|_V \|v\|_V \leq \frac{d}{dt} (S(t)u, v)_H \leq C_2 \|u\|_V \|v\|_V$ for a.e. $t \in [0, T]$ (9)

for every $u, v \in V$, for some non-negative constants C_1, C_2 and $\Gamma \in \mathbf{R}$ (observe that the derivative of $(S(t)u, v)_H$ is estimated by the norms of u and v in the space V).

Then we define an operator \mathcal{S}

$$\mathcal{S}: L^2(0, T; H) \rightarrow L^2(0, T; H) \quad \text{by} \quad \mathcal{S}u(t) := S(t)u(t) \quad (10)$$

which turns out to be linear and bounded by the constant C_1 . Since S satisfies (9) we can define a family of equibounded operators

$$S': [0, T] \rightarrow \mathcal{L}(V, V') \quad \text{by} \quad \langle S'(t)u, v \rangle_{V' \times V} := \frac{d}{dt} (S(t)u, v)_H$$

and an operator

$$S': \mathcal{V} \rightarrow \mathcal{V}' \quad \text{by} \quad \langle S'u, v \rangle_{\mathcal{V}' \times \mathcal{V}} := \int_0^T \langle S'(t)u(t), v(t) \rangle_{V' \times V} dt$$

which turns out to be linear and bounded by $\max\{|\Gamma|, C_2\}$.

Assumptions about \mathcal{N}

The only requirement about the operator \mathcal{N}

$$\mathcal{N}: \mathcal{V} \rightarrow \mathcal{V}'$$

is boundedness, i.e. there is a constant C_3 such that

$$\|\mathcal{N}u\|_{\mathcal{V}'} \leq C_3 \|u\|_{\mathcal{V}}. \quad (11)$$

Assumptions about \mathcal{A}

Consider a family of operators:

$$\begin{aligned} A(t): V \rightarrow V' \quad \text{with} \quad t \mapsto \langle A(t)u, v \rangle_{V' \times V} \text{ measurable on } [0, T], \\ A(t) \text{ linear, monotone and symmetric for a.e. } t \in [0, T], \end{aligned}$$

such that if we define the abstract operator

$$\mathcal{A}: \mathcal{V} \rightarrow \mathcal{V}', \quad \mathcal{A}u(t) = A(t)u(t), \quad 0 \leq t \leq T, \quad (12)$$

this turns out to be linear, monotone and symmetric. We moreover suppose \mathcal{A} differentiable (see Definition 2.1) and denote by \mathcal{A}' the operator

$$\mathcal{A}': \mathcal{V} \rightarrow \mathcal{V}' \quad \text{defined by} \quad \mathcal{A}'v(t) := A'(t)v(t).$$

About \mathcal{A} we suppose that there is a non-negative constant C_4 , such that

- (i) $A(t)$ symmetric for every t ,
 - (ii) $\max_{t \in [0, T]} \|A(t)\|_{\mathcal{L}(V, V')} \leq C_4$,
 - (iii) \mathcal{A} monotone,
 - (iv) $t \mapsto \langle A(t)u, v \rangle_{V' \times V}$ is absolutely continuous on $[0, T]$,
- and $\frac{d}{dt} \langle A(t)u, u \rangle_{V' \times V} \leq 0$ for every $u, v \in V$. (13)

Consider the operator

$$J: \mathcal{V} \rightarrow \mathcal{V}, \quad [Jv](t) := \int_0^t v(\sigma) d\sigma, \quad (14)$$

and a family $Q(t): V \rightarrow V'$ of linear, monotone and symmetric operators. Denote by \mathcal{Q} the operator $\mathcal{Q}: \mathcal{V} \rightarrow \mathcal{V}'$ defined by

$$\mathcal{Q}v(t) = Q(t)v(t).$$

We recall the following proposition.

Proposition 2.3. *Consider $Q(t): V \rightarrow V'$ a family of linear, monotone and symmetric operators. Suppose \mathcal{Q} differentiable and denote by $Q'(t): V \rightarrow V'$ the operators defined as follows:*

$$\langle Q'(t)w_1, w_2 \rangle_{V' \times V} := \frac{d}{dt} \langle Q(t)w_1, w_2 \rangle_{V' \times V}.$$

If $-\mathcal{Q}'(t)$ is monotone for a.e. $t \in [0, T]$ then the operator

$$\mathcal{V} \ni v \mapsto \mathcal{Q}Jv(t) = Q(t) \int_0^t v(\sigma) d\sigma$$

is monotone. If \mathcal{Q} is bounded, $\mathcal{Q}J$ is bounded by $T\|\mathcal{Q}\|_{\mathcal{L}(\mathcal{V}, \mathcal{V}')}.$

Remark 2.4. Analogously one can prove that if $Q'(t)$ is monotone for a.e. $t \in [0, T]$ and we define $J_T v(t) = \int_t^T v(\sigma) d\sigma$, then the operator $QJ: \mathcal{V} \rightarrow \mathcal{V}'$ is monotone (if Q is bounded, also QJ is bounded).

Proof. The space of polynomials $q(t)$ with coefficients in V and such that $q(0) = 0$ is dense in \mathcal{V} . Since $Q(t)$ is linear and symmetric we have

$$\frac{d}{dt} \langle Q(t)q(t), q(t) \rangle_{V' \times V} = 2 \langle Q(t)q(t), q'(t) \rangle_{V' \times V} + \langle Q'(t)q(t), q(t) \rangle_{V' \times V}. \quad (15)$$

Integrating (15) in $(0, T)$ if q is a polynomial such that $q(0) = 0$ we have that

$$2 \langle Qq, q' \rangle_{V' \times V} = - \langle Q'q, q \rangle_{V' \times V} + \langle Q(T)q(T), q(T) \rangle_{V' \times V} \geq 0$$

and taking $q(t) = Jp(t) = \int_0^t p(\sigma) d\sigma$ with p polynomial we conclude.

For the last statement observe that J is continuous and, if Q is bounded, also Q is continuous. Then QJ is linear and continuous. \square

Finally we recall the following classical result for which we refer to [16] (see Section 32.4).

Theorem 2.5. Let $B: X \rightarrow X'$ (X' the dual space of X , X Banach space) be continuous, monotone, bounded and coercive, i.e. $\lim_{\|x\| \rightarrow +\infty} \|x\|^{-1} \langle Bx, x \rangle \rightarrow +\infty$. Suppose $L: X \rightarrow X'$ to be maximal monotone. Then for every $f \in X'$ the following equation has a solution

$$Lu + Bu = f.$$

If moreover B is strictly monotone the solution is unique.

3. The existence result

In the present section we want to study the problem:

$$\begin{cases} (Su')' + \mathcal{N}u' + Au = f, \\ P_+(0)u'(0) = \varphi, \\ P_-(T)u'(T) = \psi, \\ u(0) = \eta \end{cases} \quad (16)$$

with $f \in \mathcal{V}'$, $\varphi \in \tilde{H}_+(0)$, $\psi \in \tilde{H}_-(T)$, $\eta \in V$.

The boundary conditions with respect to the variable t , i.e. the initial/final conditions, are given as follows: we give an initial condition for u' at time zero where \mathcal{S} is positive (i.e. the datum φ) while a final condition at time T where \mathcal{S} is negative (i.e. the datum ψ). Where \mathcal{S} is null, no conditions for u' are given.

About u we impose an initial datum (i.e. η) at time zero, but, as we will see, this datum could be given also at time T .

If $\mathcal{S} \equiv 0$ the initial/final conditions make no sense and the problem simply becomes

$$\begin{cases} \mathcal{N}u' + Au = f, \\ u(0) = \eta. \end{cases} \quad (17)$$

The initial/final conditions we require about u' and u are easily understood by explaining how we prove the existence result: indeed the idea to solve problem (16) is to consider the operator J defined in (14) and the change of variable $v = u'$ in (16) and then solve the first order problem:

$$\begin{cases} (Sv)' + \mathcal{N}v + \mathcal{A}Jv = f - \mathcal{A}\eta, \\ P_+(0)v(0) = \varphi, \\ P_-(T)v(T) = \psi. \end{cases} \quad (18)$$

Then it is natural to impose for the equation, partially elliptic and partially parabolic, both forward and backward, an initial datum where S is positive, no datum where $S = 0$, a final datum at time T where S is negative.

Clearly the initial condition about u is due to the choice of the operator J . In the same way one could fix a condition at time T , that is to consider $u(T) = \eta$, provided we consider the change of variable defined by the operator J_T defined in Remark 2.4.

For our purpose we introduce the space

$$\mathcal{W} = \{v \in \mathcal{V} \mid (Sv)' \in \mathcal{V}'\}, \quad \|u\|_{\mathcal{W}} = \|u\|_{\mathcal{V}} + \|(Su)'\|_{\mathcal{V}'} \quad (19)$$

and its subspace (see Proposition 2.6, Remarks 2.7 and 3.1 in [8])

$$\mathcal{W}^0 = \{v \in \mathcal{W} \mid P_+(0)v(0) = 0, P_-(T)v(T) = 0\}.$$

Definition 3.1. We say that $v \in \mathcal{W}$ is a solution of problem (18) with $f \in \mathcal{V}'$, $\varphi \in \tilde{H}_+(0)$, $\psi \in \tilde{H}_-(T)$, if

$$\begin{aligned} (Sv)'(t) + \mathcal{N}v(t) + \mathcal{A}Jv(t) &= f(t) \quad \text{in } V' \quad \text{for a.e. } t \in [0, T], \\ P_+(0)v(0) &= \varphi, \quad P_-(T)v(T) = \psi. \end{aligned}$$

We say that $u \in \{w \in C([0, T]; V) \mid w' \in \mathcal{W}\}$ is a solution of problem (16) with $f \in \mathcal{V}'$, $\varphi \in \tilde{H}_+(0)$, $\psi \in \tilde{H}_-(T)$, $u(0) = \eta$, if

$$\begin{aligned} (Su')'(t) + \mathcal{N}u'(t) + \mathcal{A}u(t) &= f(t) \quad \text{in } V' \quad \text{for a.e. } t \in [0, T], \\ P_+(0)u'(0) &= \varphi, \quad P_-(T)u'(T) = \psi, \quad u(0) = \eta. \end{aligned}$$

If $S \equiv 0$ the solution of (17) will be a function in $H^1(0, T; V)$.

Proposition 3.2. A function $v \in \mathcal{W}$ is a solution of (18) if and only if

$$\begin{aligned} -(Sv, \Phi')_{\mathcal{H}} + \langle \mathcal{N}v, \Phi \rangle_{\mathcal{V}' \times \mathcal{V}} + \langle \mathcal{A}Jv, \Phi \rangle_{\mathcal{V}' \times \mathcal{V}} \\ = \langle f, \Phi \rangle_{\mathcal{V}' \times \mathcal{V}} + (S_+(0)\varphi, \Phi(0))_H - (S_-(T)\psi, \Phi(T))_H \end{aligned} \quad (20)$$

for every $\Phi \in X = \{v \in \mathcal{V} \mid v' \in \mathcal{H}, P_-(0)v = 0, P_+(T)v = 0\}$.

Proof. Consider the definition given in Definition 3.1 and suppose v is a solution of (18). Multiplying the equation in (18) by a function $\Phi \in X$ we easily obtain (20). Suppose now to have (20) for every $\Phi \in X$. Consider

$$\Phi(t) = \vartheta(t)w + \vartheta_+(t)w_+ + \vartheta_-(t)w_-$$

where $w \in V$, $w_+ \in V \cap (\tilde{H}_+(0) \oplus \tilde{H}_0(0))$, $w_- \in V \cap (\tilde{H}_-(T) \oplus \tilde{H}_0(T))$, $\vartheta(0) = \vartheta(T) = \vartheta_+(T) = \vartheta_-(0) = 0$. Then by (20) we obtain

$$\begin{aligned} & - \int_0^T (Sv(t), \vartheta'(t)w + \vartheta'_+(t)w_+ + \vartheta'_-(t)w_-)_H dt \\ & + \int_0^T \langle \mathcal{N}v(t) + \mathcal{A}Jv(t), \vartheta(t)w + \vartheta_+(t)w_+ + \vartheta_-(t)w_- \rangle_{V' \times V} dt \\ & = \int_0^T \langle f(t), \vartheta(t)w + \vartheta_+(t)w_+ + \vartheta_-(t)w_- \rangle_{V' \times V} dt \\ & + (S_+(0)\varphi, \vartheta_+(0)w_+) - (S_-(T)\psi, \vartheta_-(T)w_-). \end{aligned} \quad (21)$$

Taking $\vartheta_+ = \vartheta_- = 0$ in (21) we obtain

$$\int_0^T \left(\frac{d}{dt} Sv(t), w \right)_H \vartheta(t) dt + \int_0^T \langle \mathcal{N}v(t) + \mathcal{A}Jv(t), w \rangle_{V' \times V} \vartheta(t) dt = \int_0^T \langle f(t), w \rangle_{V' \times V} \vartheta(t) dt.$$

Since this is true for every $\vartheta \in C_0^1(0, T)$ (and for every $w \in V$) we obtain that

$$\frac{d}{dt} (Sv(t), w)_H + \langle \mathcal{N}v(t) + \mathcal{A}Jv(t), w \rangle_{V' \times V} = \langle f(t), w \rangle_{V' \times V}$$

for every $w \in V$ and a.e. $t \in [0, T]$. From this equality, (20) and

$$\begin{aligned} & (S(T)v(T), \vartheta_-(T)w_-)_H - (S(0)v(0), \vartheta_+(0)w_+)_H \\ & = \int_0^T \frac{d}{dt} [(Sv(t), \vartheta(t)w + \vartheta_+(t)w_+ + \vartheta_-(t)w_-)_H] dt \\ & = \int_0^T \left[\frac{d}{dt} (Sv(t), w)_H \vartheta(t) + \frac{d}{dt} (Sv(t), w_+)_H \vartheta_+(t) + \frac{d}{dt} (Sv(t), w_-)_H \vartheta_-(t) \right] dt \\ & + \int_0^T [(Sv(t), w)_H \vartheta'(t) + (Sv(t), w_+)_H \vartheta'_+(t) + (Sv(t), w_-)_H \vartheta'_-(t)] dt \end{aligned}$$

we derive

$$\begin{aligned} & (S(T)v(T), \vartheta_-(T)w_-)_H - (S(0)v(0), \vartheta_+(0)w_+)_H \\ &= (S_-(T)\psi, \vartheta_-(T)w_-)_H - (S_+(0)\varphi, \vartheta_+(0)w_+)_H \end{aligned}$$

for every $\vartheta_+, \vartheta_-, w_-, w_+$ as chosen above. Then $P_+(0)v(0) = \varphi$ and $P_-(T)v(T) = \psi$. \square

Now observe that if $(Su)' \in \mathcal{V}'$ and $S'u \in \mathcal{V}'$, Su' makes sense in \mathcal{V}' . Then, if we introduce the operator

$$\mathcal{L}: D(\mathcal{L}) \subset \mathcal{V} \rightarrow \mathcal{V}', \quad \mathcal{L}u = Su' + \frac{1}{2}S'u, \quad D(\mathcal{L}) = \mathcal{W}^0,$$

the following result holds (see Proposition 3.2 in [8]).

Proposition 3.3. *The operator $\mathcal{L}: D(\mathcal{L}) \subset \mathcal{V} \rightarrow \mathcal{V}'$ is maximal monotone.*

Denote by \mathcal{B} the operator

$$\mathcal{B} = \frac{1}{2}S' + \mathcal{N} + \mathcal{A}J \quad (22)$$

and by $B(\sigma)$ the operator $B(\sigma) = \frac{1}{2}S'(\sigma) + N(\sigma) + A(\sigma)J(\sigma)$ for $\sigma \in [0, T]$, so that we write the equation $(Sv)' + \mathcal{N}v + \mathcal{A}Jv = f$ in (18) as

$$\mathcal{L}v + \mathcal{B}v = f$$

and use Theorem 2.5 to prove the result just below. At first we define the following spaces:

$$\begin{aligned} V_+(0) &= \{[P_+(0) + P_0(0)]w \in V \mid w \in V\} = V \cap (\tilde{H}_+(0) \oplus \tilde{H}_0(0)), \\ V_-(T) &= \{[P_-(T) + P_0(T)]w \in V \mid w \in V\} = V \cap (\tilde{H}_-(T) \oplus \tilde{H}_0(T)). \end{aligned}$$

Observe that the operator \mathcal{B} is bounded (by (9), (11), (13) and Proposition 2.3) by

$$\|\mathcal{B}u\|_{\mathcal{V}'} \leq C_5 \|u\|_{\mathcal{V}} \quad \text{for every } u \in \mathcal{V}, \quad C_5 = \frac{1}{2}C_2 + C_3 + TC_4. \quad (23)$$

Moreover, if we suppose there is a constant $\gamma \in \mathbf{R}$ for which (see (9) for Γ)

$$\langle \mathcal{N}u - \mathcal{N}v, u - v \rangle \geq \gamma \|u - v\|_{\mathcal{V}}^2 \quad \text{for every } u, v \in \mathcal{V}, \quad \text{and} \quad \gamma + \Gamma/2 > 0, \quad (24)$$

the operator \mathcal{B} satisfies (by Proposition 2.3)

$$\langle \mathcal{B}u - \mathcal{B}v, u - v \rangle \geq \left\langle \frac{1}{2}S'(u - v) + \mathcal{N}u - \mathcal{N}v, u - v \right\rangle \geq \left(\frac{\Gamma}{2} + \gamma \right) \|u - v\|_{\mathcal{V}}^2. \quad (25)$$

To prove the following result we will require that

$$H_+(0) \cap H_-(T) = \{0\}. \quad (26)$$

Theorem 3.4. Suppose \mathcal{B} monotone and coercive. Under assumptions (9), (11), (13), (26) for every $f \in \mathcal{V}'$, $\eta \in V$, $\varphi \in V$, $\psi \in V$ problem (16) admits one solution $u \in C([0, T]; V)$ with $u' \in \mathcal{W}$. If \mathcal{B} is strictly monotone the solution is unique. If moreover

$$V_+(0) \text{ dense in } \tilde{H}_+(0), \quad V_-(T) \text{ dense in } \tilde{H}_-(T), \quad (27)$$

\mathcal{N} is continuous and (25) holds with $\Gamma/2 + \gamma > 0$ then the data φ and ψ may be chosen respectively in $\tilde{H}_+(0)$ and $\tilde{H}_-(T)$ and the following estimate holds:

$$\begin{aligned} & \| (Su')' \|_{\mathcal{V}'} + \| u' \|_{\mathcal{V}} + \max_{t \in [0, T]} \| u(t) \|_V \\ & \leq \| \eta \|_V + c (\| f \|_{\mathcal{V}'} + \| S_-^{1/2}(T) \psi \|_{H_-(T)} + \| S_+^{1/2}(0) \varphi \|_{H_+(0)}) \end{aligned}$$

where c is a constant depending (only) on $T, C_5, \Gamma/2 + \gamma$ (see (23) and (25)).

Remark 3.5. For details about assumption (27) we refer to Section 4 in [8].

Remark 3.6. One can obtain an analogous result for the problem (16) with the condition $u(T) = \eta$ instead of $u(0) = \eta$ replacing the assumption $\frac{d}{dt} \langle A(t)u, u \rangle_{V' \times V} \leq 0$ in (13) with $\frac{d}{dt} \langle A(t)u, u \rangle_{V' \times V} \geq 0$ (see Remark 2.4).

In particular if \mathcal{A} is independent of t the problem admits a unique solution both with condition about u at time zero and at time T .

Remark 3.7. Assumption (26) means $\text{supp}(\varphi) \cap \text{supp}(\psi) = \emptyset$ (which is needed in Step 2 of the proof).

Proof. Consider the operator J defined in (14). We can rewrite problem (16) as the first-order evolution problem:

$$\begin{cases} (Sv)' + \mathcal{N}v + \mathcal{A}Jv = f - \mathcal{A}\eta, \\ P_+(0)v(0) = \varphi, \\ P_-(T)v(T) = \psi, \end{cases} \quad u(t) = \eta + \int_0^t v(\sigma) d\sigma \quad (28)$$

in the space $\mathcal{W} = \{v \in \mathcal{V} \mid (Sv)' \in \mathcal{V}'\}$. It is easy to verify that (16) and (28) are equivalent, i.e. if $u \in C([0, T]; V)$ and $u' \in \mathcal{W}$ is the solution of (16) then $v = u' \in \mathcal{W}$ is the solution of (28) and, vice versa, if $v \in \mathcal{W}$ is the solution of (28) then $u = \eta + Jv \in C([0, T]; V)$ is the solution of (16) and $u' \in \mathcal{W}$ (see, e.g., [16, Section 32.10]).

First observe that the operator \mathcal{N} is strictly monotone, bounded and coercive by (11), $\mathcal{A}J$ is continuous, monotone, bounded and coercive by (13) and Proposition 2.3. Note then that requiring

$$\mathcal{B} = \frac{1}{2}S' + \mathcal{N} + \mathcal{A}J$$

to be strictly monotone, bounded and coercive allows to \mathcal{S} to be such that

$$\langle \mathcal{S}'u, u \rangle < 0 \quad \text{for some } u \in \mathcal{V}.$$

Step 1. Suppose first $\varphi = \psi = \eta = 0$. Write the equation $(Sv)' + \mathcal{N}v + \mathcal{A}Jv = f$ as follows:

$$\left[Sv' + \frac{1}{2}S'v \right] + \left[\frac{1}{2}S'v + \mathcal{N}v + \mathcal{A}Jv \right] = f.$$

By Proposition 3.3 the operator $\mathcal{L}: \mathcal{W}^0 \rightarrow \mathcal{V}'$, $\mathcal{L}v = Sv' + \frac{1}{2}S'v$, is maximal monotone; the operator $\mathcal{B}: \mathcal{W}^0 \rightarrow \mathcal{V}'$, $\mathcal{B}v = \frac{1}{2}S'v + \mathcal{N}v + \mathcal{A}Jv$ is monotone, bounded and coercive by assumption. Then we can use Theorem 2.5 to conclude that

$$\begin{cases} (Sv)' + \mathcal{N}v + \mathcal{A}Jv = f, \\ P_+(0)v(0) = 0, \\ P_-(T)v(T) = 0 \end{cases}$$

has one solution. If \mathcal{B} is strictly monotone the solution is unique.

Step 2. Suppose \mathcal{B} is strictly monotone and consider first $\Phi, \Psi \in V$ with $P_+(0)\Phi = \varphi$, $P_-(T)\Psi = \psi$, $P_+(0)\Psi = 0$, $P_-(T)\Phi = 0$ (and still $\eta = 0$). This is possible because of (26). We first define $\vartheta = \Phi + \Psi$ and solve

$$\begin{cases} (Sw)' + \mathcal{N}w + \mathcal{A}Jw = f - S'\vartheta - \mathcal{N}\vartheta - \mathcal{A}J\vartheta, \\ P_+(0)w(0) = 0, \\ P_-(T)w(T) = 0. \end{cases}$$

We can solve this problem by Step 1. Then, denoting by w a solution, $v = w + \vartheta$ will be a solution of the first order equation in (28) since, thanks to (26), $P_+(0)\vartheta = P_+(0)\Phi$ and $P_-(T)\vartheta = P_-(T)\Psi$.

Step 3. Suppose now \mathcal{N} is continuous, (24) and (27) to be true and consider $\varphi \in \tilde{H}_+(0)$ and $\psi \in \tilde{H}_-(T)$. Consider two sequences $(\Phi_n)_n \subset V_+(0)$ and $(\Psi_n)_n \subset V_-(T)$ such that (thanks to (27))

$$\varphi_n := P_+(0)\Phi_n \rightarrow \varphi \quad \text{in } \tilde{H}_+(0), \quad \psi_n := P_-(T)\Psi_n \rightarrow \psi \quad \text{in } \tilde{H}_-(T),$$

$f \in \mathcal{V}'$, $\eta = 0$ and the problems:

$$\begin{cases} (Su')' + \mathcal{N}u' + \mathcal{A}u = f, \\ P_+(0)u'(0) = \varphi_n, \\ P_-(T)u'(T) = \psi_n, \\ u(0) = 0. \end{cases}$$

Denote by v_n the solutions (by Step 2) of the problems:

$$\begin{cases} (Sv)' + \mathcal{N}v + \mathcal{A}Jv = f, \\ P_+(0)v(0) = \varphi_n, \\ P_-(T)v(T) = \psi_n. \end{cases}$$

Consider the $v_n - v_k$: since the problems above are linear the function $v_n - v_k$ solves the problem:

$$\begin{cases} (\mathcal{S}v)' + \mathcal{N}v + \mathcal{A}Jv = 0, \\ P_+(0)v(0) = \varphi_n - \varphi_k, \\ P_-(T)v(T) = \psi_n - \psi_k. \end{cases}$$

Multiplying the equation by $v_n - v_k$ we obtain

$$\begin{aligned} & \left\langle \left[\frac{1}{2} \mathcal{S}' + \mathcal{N} + \mathcal{A}J \right] (v_n - v_k), v_n - v_k \right\rangle_{\mathcal{V}' \times \mathcal{V}} \\ &= - \left\langle \mathcal{S}(v_n - v_k)' + \frac{1}{2} \mathcal{S}'(v_n - v_k), v_n - v_k \right\rangle_{\mathcal{V}' \times \mathcal{V}} \\ &= \frac{1}{2} \left[(\mathcal{S}(0)(v_n - v_k)(0), (v_n - v_k)(0))_H - (\mathcal{S}(T)(v_n - v_k)(T), (v_n - v_k)(T))_H \right] \\ &\leq \frac{1}{2} \left[(\mathcal{S}_+(0)(v_n - v_k)(0), (v_n - v_k)(0))_H + (\mathcal{S}_-(T)(v_n - v_k)(T), (v_n - v_k)(T))_H \right]. \end{aligned}$$

By assumption this term is going to zero for $n, k \rightarrow +\infty$. In particular, by assumption, $\gamma + \Gamma/2 > 0$ and by (25) we have

$$\begin{aligned} (\gamma + \Gamma/2) \|v_n - v_k\|_{\mathcal{V}}^2 &\leq \left\langle \left[\frac{1}{2} \mathcal{S}' + \mathcal{N} \right] (v_n - v_k), v_n - v_k \right\rangle_{\mathcal{V}' \times \mathcal{V}} \\ &\leq \left\langle \left[\frac{1}{2} \mathcal{S}' + \mathcal{N} + \mathcal{A}J \right] (v_n - v_k), v_n - v_k \right\rangle_{\mathcal{V}' \times \mathcal{V}}. \end{aligned}$$

Then $\{v_n\}_n$ is a Cauchy sequence in \mathcal{V} and consequently

$$v_n \rightarrow v \quad \text{and} \quad Jv_n \rightarrow Jv \quad \text{in } \mathcal{V}.$$

By Proposition 3.2 we get

$$\begin{aligned} & -(\mathcal{S}v_n, \Phi')_{\mathcal{H}} + (\mathcal{N}v_n, \Phi)_{\mathcal{V}' \times \mathcal{V}} + (\mathcal{A}Jv_n, \Phi)_{\mathcal{V}' \times \mathcal{V}} \\ &= (f, \Phi)_{\mathcal{V}' \times \mathcal{V}} + (\mathcal{S}_+(0)\varphi_n, \Phi(0))_H - (\mathcal{S}_-(T)\psi_n, \Phi(T))_H \end{aligned}$$

for every $\Phi \in \{v \in \mathcal{V} \mid v' \in \mathcal{H}, P_-(0)v = 0, P_+(T)v = 0\}$. Taking the limit and by the continuity of \mathcal{S} , \mathcal{N} and $\mathcal{A}J$ we obtain a limit equation satisfied by the function v with $\varphi \in \tilde{H}_+(0)$ and $\psi \in \tilde{H}_-(T)$.

Step 4. Finally, if $\eta \in V$ is not null, we consider u the solution of the problem:

$$\begin{cases} (\mathcal{S}u)' + \mathcal{N}u' + \mathcal{A}u = f - \mathcal{A}\eta, \\ P_+(0)u'(0) = \varphi, \\ P_-(T)u'(T) = \psi, \\ u(0) = 0 \end{cases}$$

with $f \in \mathcal{V}'$, $\varphi \in \tilde{H}_+(0)$, $\psi \in \tilde{H}_-(T)$, and then consider $\eta + u(t)$, which solves (16).

To conclude and prove the estimate observe that, by Theorem 3.8 in [8], we have that

$$\|(\mathcal{S}u')'\|_{\mathcal{V}'} + \|u'\|_{\mathcal{V}} \leq c(\|f\|_{\mathcal{V}'} + \|S_-^{1/2}(T)\psi\|_{H_-(T)} + \|S_+^{1/2}(0)\varphi\|_{H_+(0)})$$

where $c = c(C_5, \Gamma/2 + \gamma)$. Finally we conclude by

$$\|u(t)\|_V = \left\| \eta + \int_0^t u'(\sigma) d\sigma \right\|_V \leq \|\eta\|_V + T^{1/2} \|u'\|_{\mathcal{V}}. \quad \square$$

Here we present a concrete example, just to explain the assumptions about the operators \mathcal{S} , \mathcal{N} , \mathcal{A} , for further examples of possible choices of the function s in (29) we refer to [8].

Consider $T > 0$, Ω open bounded subset of \mathbf{R}^n with Lipschitzian boundary and a function

$$s : \Omega \times [0, T] \rightarrow \mathbf{R}.$$

By s_+ and s_- we denote respectively the non-negative and non-positive part of s . Consider the following problem:

$$\begin{cases} (su_t)_t - \operatorname{div}(\alpha(|Du_t|^2)Du_t) + ru_t - \operatorname{div}(a \cdot Du) = f(x, t) & \text{on } \Omega \times (0, T), \\ u(x, t) = 0, & (x, t) \in \partial\Omega \times (0, T), \\ \int_{\Omega} (u_t(x, 0) - \varphi(x))^2 s_+(x, 0) dx = 0, \\ \int_{\Omega} (u_t(x, T) - \psi(x))^2 s_-(x, T) dx = 0, \\ u(x, 0) = \eta(x), & x \in \Omega. \end{cases} \quad (29)$$

The initial/final conditions $P_+(0)u'(0) = \varphi$ and $P_-(T)u'(T) = \psi$ in (16) are to be understood as the integral conditions involving φ and ψ in (29).

Notice that if $s \equiv 1$, $r \equiv 0$ and $a_{ij} = \delta_{ij}$ this equation is the wave equation with a dissipative term and with initial condition φ for u_t on the whole Ω (the condition involving ψ disappears).

This problem can be thought of as a model problem for (16): consider $V = H_0^1(\Omega)$, $H = L^2(\Omega)$, $\mathcal{V} = L^2(0, T; H_0^1(\Omega))$, $\mathcal{H} = L^2(0, T; L^2(\Omega)) = L^2(\Omega \times (0, T))$ and

$$S(t) : H \rightarrow H \quad \text{defined by} \quad S(t)u = s(x, t)u(x)$$

and then

$$\mathcal{S} : \mathcal{H} \rightarrow \mathcal{H} \quad \text{defined by} \quad \mathcal{S}u(t)(x) = s(x, t)u(x, t). \quad (30)$$

In order that (9) is satisfied, the function s should be such that

$$t \mapsto \int_{\Omega} u(x)v(x)s(x, t) dx \quad \text{is absolutely continuous,}$$

and for this reason we can consider the integral conditions involving φ and ψ in (29), and

$$\Gamma \int_{\Omega} |Du(x)|^2 dx \leq \frac{d}{dt} \int_{\Omega} u^2(x) s(x, t) dx \leq C_2 \int_{\Omega} |Du(x)|^2 dx$$

with $\Gamma \in \mathbf{R}$, $C_2 \geq 0$ (for more details we refer to [8]). The data will satisfy: $f \in L^2(0, T; H^{-1}(\Omega))$, $\varphi \in L^2(\Omega_+(0), s_+(\cdot, 0))$, $\psi \in L^2(\Omega_-(T), s_-(\cdot, T))$, $\eta \in H_0^1(\Omega)$ (for the choice of φ and ψ in such spaces we refer to [8]). The operator \mathcal{N} is defined by

$$\mathcal{N}: \mathcal{V} \rightarrow \mathcal{V}', \quad \mathcal{N}w(t)(x) = -\operatorname{div}(\alpha(|Dw(x, t)|^2) Dw(x, t))$$

with α satisfying

$$\alpha: [0, +\infty) \rightarrow [0, C], \quad \alpha(\tau^2)\tau - \alpha(\sigma^2)\sigma \geq \gamma(\tau - \sigma)$$

where C is a positive constant, γ is the constant appearing in (24) and $0 \leq \sigma < \tau$. The function $r = r(x)$ belongs to $L^\infty(\Omega)$ and finally

$$\mathcal{A}: \mathcal{V} \rightarrow \mathcal{V}', \quad \mathcal{A}w(t)(x) = -\operatorname{div}(a(x, t) \cdot Dw(x, t))$$

with $a = [a_{ij}(x, t)]_{i,j=1}^n$ satisfying (so that assumptions of Proposition 2.3 is satisfied)

$$a_{ij} = a_{ji}, \quad \int_0^T \int_{\Omega} \sum_{i,j=1}^n (a_{ij})_t D_i u D_j u dx dt \leq 0.$$

The last condition is satisfied, for instance, if $a_{ij} = a_{ij}(x)$. The simpler situation could be the following: if

$$\alpha(\tau) \equiv \gamma \in \mathbf{R}, \quad r \equiv 0, \quad a_{ij} = \delta_{ij}$$

problem (29) becomes

$$\begin{cases} (su_t)_t - \gamma \Delta u_t - \Delta u = f(x, t) & \text{on } \Omega \times (0, T), \\ u(x, t) = 0, & (x, t) \in \partial\Omega \times (0, T), \\ \int_{\Omega} (u_t(x, 0) - \varphi(x))^2 s_+(x, 0) dx = 0, \\ \int_{\Omega} (u_t(x, T) - \psi(x))^2 s_-(x, T) dx = 0, \\ u(x, 0) = \eta(x), & x \in \Omega. \end{cases} \quad (31)$$

In this situation we may replace the last initial condition by $u(x, T) = \eta(x)$ in Ω , and it has a unique solution (see Remark 3.6).

We conclude by observing that if the constant Γ in (9)(iv) is strictly positive we can admit also the function α not to be non-negative. For instance, consider for simplicity example (31), we can admit also $\gamma \leq 0$ and consider in particular the equation

$$(su_t)_t - \Delta u = f$$

(where indeed, if s is regular enough, we have $su_{tt} + s_t u_t - \Delta u = f$, so a term involving u_t appears in any case).

Acknowledgment

The author thanks the referee for her/his useful remarks.

References

- [1] M.S. Baouendi, P. Grisvard, Sur une équation d'évolution changeant de type, *J. Funct. Anal.* 2 (1968) 352–367.
- [2] R.W. Carroll, R.E. Showalter, *Singular and Degenerate Cauchy Problems*, Academic Press, New York, 1976.
- [3] B.L. Keyfitz, M. Shearer (Eds.), *Nonlinear Evolution Equations that Change Type*, in: *IMA Vol. Math. Appl.*, vol. 27, Springer, New York, 1999.
- [4] L.D. Landau, E.M. Lifshitz, *Theory of Elasticity*, Pergamon Press/Addison Wesley, London–Paris–Frankfurt/Reading, MA, 1970.
- [5] J.L. Lions, E. Magenes, *Problèmes aux limites non homogènes et applications*, vol. 1, Dunod, Paris, 1968.
- [6] C.S. Morawetz, Mixed equations and transonic flow, *J. Hyperbolic Differential Equations* 1 (1) (2004) 1–26.
- [7] C.D. Pagani, G. Talenti, On a forward–backward parabolic equation, *Ann. Mat. Pura Appl.* 90 (1971) 1–57.
- [8] F. Paronetto, Existence results for a class of evolution equations of mixed type, *J. Funct. Anal.* 212 (2) (2004) 324–356.
- [9] R. Pego, Phase transition in one-dimensional nonlinear viscoelasticity: Admissibility and stability, *Arch. Ration. Mech. Anal.* 97 (1987) 353–394.
- [10] M. Potier-Ferry, On the mathematical foundations of elastic stability theory I, *Arch. Ration. Mech. Anal.* 78 (1) (1982) 55–72.
- [11] P. Rybka, Dynamical model of phase transition by means of viscoelasticity in many dimensions, *Proc. Roy. Soc. Edinburgh Sect. A* 121 (1–2) (1992) 101–138.
- [12] P. Rybka, K.H. Hoffmann, Convergence of solutions to the equation of quasi-static approximation of viscoelasticity with capillarity, *J. Math. Anal. Appl.* 226 (1998) 61–81.
- [13] R.E. Showalter, *Hilbert Space Methods for Partial Differential Equations*, Electron. Monogr. Differ. Equ., Texas State Univ., San Marcos, TX, 1994, iii + 242 pp. Available at <http://ejde.math.swt.edu/>.
- [14] R.E. Showalter, *Monotone Operators in Banach Space and Nonlinear Partial Differential Equations*, Amer. Math. Soc., 1997.
- [15] P.J. Swart, P.J. Holmes, Energy minimization and the formation of microstructure in dynamics anti-plane shear, *Arch. Ration. Mech. Anal.* 121 (1992) 37–85.
- [16] E. Zeidler, *Nonlinear Functional Analysis and Its Applications*, vols. II A and II B, Springer, New York, 1990.